

Renormalization Group Flow and the Dark Energy Problem

Shoichi Ichinose*

January 11, 2011

*Laboratory of Physics, School of Food and Nutritional Sciences, University of Shizuoka
Yada 52-1, Shizuoka 422-8526, Japan*

Abstract

Casimir energy is calculated for 5D scalar theory in the *warped* geometry. A new regularization, called *sphere lattice regularization*, is taken. The regularized configuration is *closed-string like*. We numerically evaluate Λ (4D UV-cutoff), ω (5D bulk curvature, extra space UV-boundary parameter) and T (extra space IR-boundary parameter) dependence of Casimir energy. 5D Casimir energy is *finitely* obtained after the *proper renormalization procedure*. The *warp parameter* ω suffers from the *renormalization effect*. Regarding Casimir energy as the main contribution to the cosmological term, we examine the dark energy problem.

1. Introduction In the quest for the unified theory, the higher dimensional (HD) approach is a fascinating one from the geometrical point. Historically the initial successful one is the Kaluza-Klein model, which unifies the photon, graviton and dilaton from the 5D space-time approach. The HD theories, however, generally have the serious defect as the quantum field theory(QFT) : un-renormalizability. The HD quantum field theories, at present, are not defined within the QFT.

In 1983, the Casimir energy in the Kaluza-Klein theory was calculated by Appelquist and Chodos[1]. They took the cut-off (Λ) regularization and found the quintic (Λ^5) divergence and the finite term. The divergent term shows the *unrenormalizability* of the 5D theory, but the finite term looks meaningful[2] and, in fact, is widely regarded as the right vacuum energy which shows *contraction* of the extra axis.

In the development of the string and D-brane theories, a new approach to the renormalization group was found. It is called *holographic renormalization*. We regard the renormalization group flow as a curve in the bulk. The flow goes along the extra axis. The curve is derived as a dynamical equation such as Hamilton-Jacobi equation. It originated

*Email address: ichinose@u-shizuoka-ken.ac.jp

from the AdS/CFT correspondence. Spiritually the present basic idea overlaps with this approach.

2. Casimir Energy of 5D Scalar Theory In the warped geometry, $ds^2 = \frac{1}{\omega^2 z^2}(\eta_{\mu\nu} dx^\mu dx^\nu + dz^2)$, we consider the 5D massive *scalar* theory with $m^2 = -4\omega^2 (< 0)$. $\mathcal{L} = \sqrt{-G}(-\frac{1}{2}\nabla^A \Phi \nabla_A \Phi - \frac{1}{2}m^2 \Phi^2)$. The Casimir energy E_{Cas} is given by

$$e^{-T^{-4}E_{Cas}} = \int \mathcal{D}\Phi \exp\left\{i \int d^5 X \mathcal{L}\right\} \Big|_{\text{Euclid}} = \exp \sum_{n,p} \left\{-\frac{1}{2} \ln(p_E^2 + M_n^2)\right\}, \quad p_E^2 \equiv p_1^2 + p_2^2 + p_3^2 + p_4^2, \quad (1)$$

where M_n is the eigenvalues of the following operator.

$$\{s(z)^{-1} \hat{L}_z + M_n^2\} \psi_n(z) = 0 \quad , \quad \hat{L}_z \equiv \frac{d}{dz} \frac{1}{(\omega z)^3} \frac{d}{dz} - \frac{m^2}{(\omega z)^5} \quad , \quad (2)$$

where $s(z) = \frac{1}{(\omega z)^3}$. Z_2 parity is imposed as: $\psi_n(z) = -\psi_n(-z)$ for $P = -$; $\psi_n(z) = \psi_n(-z)$ for $P = +$. The expression (1) is the familiar one of the Casimir energy. It is re-expressed in a *closed* form using the heat-kernel method and the propagator as follows. First we can express it, using the heat equation solution, as ($\omega/T = e^{\omega l}$, l is the periodicity in y -coordinate: $y \rightarrow y + 2l, \omega|z| = e^{\omega|y|}$),

$$e^{-T^{-4}E_{Cas}} = (\text{const}) \times \exp \left[T^{-4} \int \frac{d^4 p}{(2\pi)^4} 2 \int_0^\infty \frac{1}{2} \frac{dt}{t} \text{Tr} H_p(z, z'; t) \right] \quad ,$$

$$\text{Tr} H_p(z, z'; t) = \int_{1/\omega}^{1/T} s(z) H_p(z, z'; t) dz \quad , \quad \left\{ \frac{\partial}{\partial t} - (s^{-1} \hat{L}_z - p^2) \right\} H_p(z, z'; t) = 0 \quad . \quad (3)$$

The heat kernel $H_p(z, z'; t)$ is formally solved, using the Dirac's bra and ket vectors ($z|, |z$), as $H_p(z, z'; t) = (z| e^{-(s^{-1} \hat{L}_z + p^2)t} |z')$. We here introduce the position/momentum propagators $G_p^\mp$: $G_p^\mp(z, z') \equiv \int_0^\infty dt H_p(z, z'; t)$. They satisfy the following differential equations of *propagators*.

$$(\hat{L}_z - p^2 s(z)) G_p^\mp(z, z') = \begin{cases} \epsilon(z) \epsilon(z') \hat{\delta}(|z| - |z'|) & P = -1 \\ \hat{\delta}(|z| - |z'|) & P = 1 \end{cases} \quad , \quad p^2 = p_\mu p^\mu = -p_0^2 + p_1^2 + p_2^2 + p_3^2 \quad (4)$$


$G_p^\mp$ can be expressed in a *closed* form. Taking the *Dirichlet* condition at all fixed points, the expression for the fundamental region ($1/\omega \leq z \leq z' \leq 1/T$) is given by

$$G_p^\mp(z, z') = \mp \frac{\omega^3}{2} z^2 z'^2 \frac{\{\mathbf{I}_0(\frac{\tilde{p}}{\omega}) \mathbf{K}_0(\tilde{p}z) \mp \mathbf{K}_0(\frac{\tilde{p}}{\omega}) \mathbf{I}_0(\tilde{p}z)\} \{\mathbf{I}_0(\frac{\tilde{p}}{T}) \mathbf{K}_0(\tilde{p}z') \mp \mathbf{K}_0(\frac{\tilde{p}}{T}) \mathbf{I}_0(\tilde{p}z')\}}{\mathbf{I}_0(\frac{\tilde{p}}{T}) \mathbf{K}_0(\frac{\tilde{p}}{\omega}) - \mathbf{K}_0(\frac{\tilde{p}}{T}) \mathbf{I}_0(\frac{\tilde{p}}{\omega})} \quad , \quad (5)$$

where $\tilde{p} \equiv \sqrt{p^2}$, $p^2 \geq 0$. I_0 and K_0 are modified Bessel functions. We can express Casimir energy as,

$$-E_{Cas}^{\Lambda, \mp}(\omega, T) = \int \frac{d^4 p_E}{(2\pi)^4} \Big|_{\tilde{p} \leq \Lambda} \int_{1/\omega}^{1/T} dz F^\mp(\tilde{p}, z), F^\mp(\tilde{p}, z) = \frac{2}{(\omega z)^3} \int_{\tilde{p}}^\Lambda \tilde{k} G_k^\mp(z, z) d\tilde{k} \quad , \quad (6)$$

Figure 1: Rectangle region of $(z, \tilde{p})$ for the integration (6). The hyperbolic curve was proposed[3].




where $\tilde{p} = \sqrt{p_E^2}$. The momentum symbol p_E indicates Euclideanization. Here we introduce the UV cut-off parameter Λ in the 4D momentum space.

3. UV and IR Regularization and Evaluation of Casimir Energy The integral region of the above equation (6) is displayed, in Fig.1, as a rectangle. In the figure, we introduce the regularization cut-offs for the 4D-momentum integral, $\mu \leq \tilde{p} \leq \Lambda$. For simplicity, we take the following IR cutoff of 4D momentum : $\mu = \Lambda \cdot \frac{T}{\omega} = \Lambda e^{-\omega l}$. Importantly, (6) shows the *scaling* behavior for large values of Λ and $1/T$. From a *close* numerical analysis, we have confirmed : (4A) $E_{Cas}^{\Lambda,-}(\omega, T) = \frac{2\pi^2}{(2\pi)^4} \times \left[-0.0250 \frac{\Lambda^5}{T} \right]$. The Λ^5 -divergence, (4A), shows the notorious problem of the higher dimensional theories. We have proposed an approach to solve this problem and given a legitimate explanation within the 5D QFT[4, 5]. See Fig.2. The IR and UV cutoffs *change* along the *etra* axis. Their S^3 -radii are given by $r_{IR}(z) = 1/\tilde{p}_{IR}(z)$ and $r_{UV}(z) = 1/\tilde{p}_{UV}(z)$. The 5D volume region bounded by B_{UV} and B_{IR} is the integral region of the Casimir energy E_{Cas} . (We call this regularization *sphere lattice regularization* because one big 4D-ball (radius r_{IR}) are made of many small 4D-balls (cells, radius r_{UV}). See Fig.12 of Ref.[6]) The forms of $r_{UV}(z)$ and $r_{IR}(z)$ can be determined by the *minimal area principle*: $3 + \frac{4}{z}r'r - \frac{r''r}{r'^2+1} = 0$, $r' \equiv \frac{dr}{dz}$, $r'' \equiv \frac{d^2r}{dz^2}$, $1/\omega \leq z \leq 1/T$. We have confirmed, by *numerically* solving the above differential equation (Runge-Kutta), those curves that show the flow of renormalization really appear. The results imply the *boundary conditions* determine the property of the renormalization flow.

4. Weight Function and the Meaning We consider another approach which respects the *minimal area principle*. Let us introduce, instead of restricting the integral

Figure 2: Region of $(\tilde{p}, z)$ for the integration (present proposal).



region, a *weight function* $W(\tilde{p}, z)$ in the $(\tilde{p}, z)$ -space for the purpose of *suppressing* UV and IR divergences of the Casimir Energy.

$$\begin{aligned}
 -E_{Cas}^{\mp W}(\omega, T) &\equiv \int \frac{d^4 p_E}{(2\pi)^4} \int_{1/\omega}^{1/T} dz W(\tilde{p}, z) F^{\mp}(\tilde{p}, z) \quad , \quad \tilde{p} = \sqrt{p_4^2 + p_1^2 + p_2^2 + p_3^2} \quad , \\
 \left\{ \begin{array}{ll} (N_1)^{-1} e^{-(1/2)\tilde{p}^2/\omega^2 - (1/2)z^2 T^2} \equiv W_1(\tilde{p}, z), \quad N_1 = 1.711/8\pi^2 & \text{elliptic suppr.} \\ (N_2)^{-1} e^{-\tilde{p}zT/\omega} \equiv W_2(\tilde{p}, z), \quad N_2 = 2\frac{\omega^3}{T^3}/8\pi^2 & \text{hyperbolic suppr.1} \\ (N_8)^{-1} e^{-1/2(\tilde{p}^2/\omega^2 + 1/z^2 T^2)} \equiv W_8(\tilde{p}, z), \quad N_8 = 0.4177/8\pi^2 & \text{reciprocal suppr.1} \end{array} \right. \quad (7)
 \end{aligned}$$

where $F^{\mp}(\tilde{p}, z)$ are defined in (6). They (except W_2) give, after normalizing the factor Λ/T , *only* the *log-divergence*.

$$E_{Cas}^W/\Lambda T^{-1} = -\alpha\omega^4 (1 - 4c \ln(\Lambda/\omega) - 4c' \ln(\Lambda/T)) \quad , \quad (8)$$

where the numerical values of α, c and c' are obtained depending on the choice of the weight function[6]. This means the 5D Casimir energy is *finitely* obtained by the ordinary renormalization of the warp factor ω . (See the final section.)

In the previous work[5], we have presented the following idea to define the weight function $W(\tilde{p}, z)$. In the evaluation (7), the $(\tilde{p}, z)$ -integral is over the rectangle region shown in Fig.1 (with $\Lambda \rightarrow \infty$ and $\mu \rightarrow 0$). Following Feynman[7], we can replace the integral by the *summation* over *all possible pathes* $\tilde{p}(z)$.

$$-E_{Cas}^W(\omega, T) = \int \mathcal{D}\tilde{p}(z) \int_{1/\omega}^{1/T} dz S[\tilde{p}(z), z], S[\tilde{p}(z), z] = \frac{2\pi^2}{(2\pi)^4} \tilde{p}(z)^3 W(\tilde{p}(z), z) F^{\mp}(\tilde{p}(z), z). \quad (9)$$

There exists the *dominant path* $\tilde{p}_W(z)$ which is determined by the minimal principle : $\delta S = 0$. Dominant Path $\tilde{p}_W(z)$: $\frac{d\tilde{p}}{dz} = -\frac{\partial \ln(WF)}{\partial z} / (\frac{3}{\tilde{p}} + \frac{\partial \ln(WF)}{\partial \tilde{p}})$. Hence it is fixed by the weight function $W(\tilde{p}, z)$. On the other hand, there exists another independent path: the minimal surface curve $r_g(z)$. Minimal Surface Curve $r_g(z)$: $3 + \frac{4}{z}r'r - \frac{r''r}{r'^2+1} = 0$, $\frac{1}{\omega} \leq z \leq \frac{1}{T}$. It is obtained by the *minimal area principle*: $\delta A = 0$ where

$$ds^2 = (\delta_{ab} + \frac{x^a x^b}{(rr')^2}) \frac{dx^a dx^b}{\omega^2 z^2} \equiv g_{ab}(x) dx^a dx^b, A = \int \sqrt{g} d^4x = \int_{1/\omega}^{1/T} \frac{\sqrt{r'^2+1} r^3}{\omega^4 z^4} dz. \quad (10)$$

Hence $r_g(z)$ is fixed by the *induced geometry* $g_{ab}(x)$. Here we put the *requirement*[5]: (4A) $\tilde{p}_W(z) = \tilde{p}_g(z)$, where $\tilde{p}_g \equiv 1/r_g$. This means the following things. We *require* the dominant path coincides with the minimal surface line $\tilde{p}_g(z) = 1/r_g(z)$ which is defined independently of $W(\tilde{p}, z)$. $W(\tilde{p}, z)$ is defined here by the induced geometry $g_{ab}(x)$. In this way, we can connect the integral-measure over the 5D-space with the geometry. We have confirmed the coincidence by the numerical method.

In order to most naturally accomplish the above requirement, we can go to a *new step*. Namely, we *propose* to *replace* the 5D space integral with the weight W , (7), by the following *path-integral*. We *newly define* the Casimir energy in the higher-dimensional theory as follows.

$$- \mathcal{E}_{Cas}(\omega, T, \Lambda) = \int_{1/\Lambda}^{1/\mu} d\rho \int \frac{r(\omega^{-1})}{= r(T^{-1})} \prod_{a,z} \mathcal{D}x^a(z) F(\frac{1}{r}, z) \exp \left[- \int_{1/\omega}^{1/T} \frac{\sqrt{r'^2+1} r^3}{2\alpha' \omega^4 z^4} dz \right], \quad (11)$$

where $\mu = \Lambda T / \omega$ and the limit $\Lambda T^{-1} \rightarrow \infty$ is taken. The string (surface) tension parameter $1/2\alpha'$ is introduced. (Note: Dimension of α' is [Length]⁴.) $F(\tilde{p}, z)$ is defined in (6) and represents the contribution from the *field-quantization* of the bulk scalar (or EM) fields. (This proposal is shown to be valid in arXiv:1004.2573, 1010.5558.)

5. Discussion and Conclusion When c and c' in (8) are sufficiently small we find the renormalization group function for the warp factor ω as

$$\omega_r = \omega(1 - c \ln(\Lambda/\omega) - c' \ln(\Lambda/T)) , \quad \beta \equiv \frac{\partial}{\partial(\ln \Lambda)} \ln \frac{\omega_r}{\omega} = -c - c' . \quad (12)$$

No local counterterms are necessary.

Through the Casimir energy calculation, in the higher dimension, we find a way to quantize the higher dimensional theories within the QFT framework. The quantization *with respect to the fields* (except the gravitational fields $G_{AB}(X)$) is done in the standard way. After this step, the expression has the summation *over the 5D space(-time) coordinates or momenta* $\int dz \prod_a dp^a$. We have proposed that this summation should be replaced by the *path-integral* $\int \prod_{a,z} \mathcal{D}p^a(z)$ with the *area action* (Hamiltonian) $A = \int \sqrt{\det g_{ab}} d^4x$ where g_{ab} is the *induced metric* on the 4D surface. This procedure says the 4D momenta p^a (or coordinates x^a) are *quantum statistical operators* and the extra-coordinate z is the inverse temperature (Euclidean time). We recall the similar situation occurs in the standard string approach. The space-time coordinates obey some uncertainty principle[8].

Recently the dark energy (as well as the dark matter) in the universe is a hot subject. It is well-known that the dominant candidate is the cosmological term. The cosmological constant λ appears as: (5A) $R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R - \lambda g_{\mu\nu} = T_{\mu\nu}^{matter}$, $S = \int d^4x \sqrt{-g} \{ \frac{1}{G_N} (R + \lambda) \} + \int d^4x \sqrt{-g} \{ \mathcal{L}_{matter} \}$, $g = \det g_{\mu\nu}$. We consider here the 3+1 dim Lorentzian space-time ($\mu, \nu = 0, 1, 2, 3$). The constant λ observationally takes the value : (5B) $\frac{1}{G_N} \lambda_{obs} \sim \frac{1}{G_N R_{cos}^2} \sim m_\nu^4 \sim (10^{-3} eV)^4$, $\lambda_{obs} \sim \frac{1}{R_{cos}^2} \sim 4 \times 10^{-66} (eV)^2$, where $R_{cos} \sim 5 \times 10^{32} eV^{-1}$ is the cosmological size (Hubble length), m_ν is the neutrino mass. On the other hand, we have theoretically so far : (5C) $\frac{1}{G_N} \lambda_{th} \sim \frac{1}{G_N^2} = M_{pl}^4 \sim (10^{28} eV)^4$. We have the famous huge discrepancy factor : (5D) $\frac{\lambda_{th}}{\lambda_{obs}} \sim N_{DL}^2$, $N_{DL} \equiv M_{pl} R_{cos} \sim 6 \times 10^{60}$, where N_{DL} is here introduced as the Dirac's large number. If we use the present result, we can obtain a natural choice of T, ω and Λ as follows. By identifying $T^{-4} E_{Cas} = -\alpha_1 \Lambda T^{-1} \omega^4 / T^4$ with $\int d^4x \sqrt{-g} (1/G_N) \lambda_{ob} = R_{cos}^2 (1/G_N)$, we obtain the following relation: (5E) $N_{DL}^2 = R_{cos}^2 \frac{1}{G_N} = -\alpha_1 \frac{\omega^4 \Lambda}{T^5}$. The warped (AdS₅) model predicts the cosmological constant *negative*, hence we have interest only in its absolute value. We take the following choice for Λ and ω : (5F) $\Lambda = M_{pl} \sim 10^{19} GeV$, $\omega \sim \frac{1}{\sqrt[4]{G_N R_{cos}^2}} = \sqrt{\frac{M_{pl}}{R_{cos}}} \sim m_\nu \sim 10^{-3} eV$.

As shown above, we have the standpoint that the cosmological constant is mainly made from the Casimir energy. We do not yet succeed in obtaining the value α_1 negatively, but succeed in obtaining the finiteness of the cosmological constant and its gross absolute value. The smallness of the value is naturally explained by the renormalization group flow. Because we already know the warp parameter ω *flows* (12), the $\lambda_{obs} \sim 1/R_{cos}^2 \propto \omega^4$, says that the *smallness of the cosmological constant comes from the renormalization group flow* for the non asymptotic-free case ($c + c' < 0$ in (12)).

The IR parameter T , the normalization factor Λ/T in (8) and the IR cutoff $\mu = \Lambda \frac{T}{\omega}$ are given by : (5G) $T = R_{cos}^{-1} (N_{DL})^{1/5} \sim 10^{-20} eV$, $\frac{\Lambda}{T} = (N_{DL})^{4/5} \sim 10^{50}$, $\mu = M_{pl} N_{DL}^{-3/10} \sim 1 GeV \sim m_N$, where m_N is the nucleon mass. The degree of freedom of the universe (space-time) is given by : (5H) $\frac{\Lambda^4}{\mu^4} = \frac{\omega^4}{T^4} = N_{DL}^{6/5} \sim 10^{74} \sim (\frac{M_{pl}}{m_N})^4$.

References

- [1] T. Appelquist and A. Chodos, Phys.Rev.Lett.**50**(1983)141
- [2] S. Ichinose, Phys.Lett.**152B**(1985),56
- [3] L. Randall and M.D. Schwartz, *JHEP* **0111** (2001) 003, hep-th/0108114
- [4] S. Ichinose and A. Murayama, Phys.Rev.**D76**(2007)065008, hep-th/0703228
- [5] S. Ichinose, Prog.Theor.Phys.**121**(2009)727, arXiv:0801.3064v8[hep-th]
- [6] S. Ichinose, arXiv:0812.1263[hep-th]
- [7] R.P. Feynman, *Statistical Mechanics*, W.A.Benjamin,Inc., Massachusetts, 1972

[8] T. Yoneya, Prog.Theor.Phys.**103**(2000)1081